

Scattering rate, transport and specific heat in a metal close to a quantum critical point : emergence of a robust Fermi liquid picture ?

George Kastinakis

Institute of Electronic Structure and Laser (IESL),

*Foundation for Research and Technology - Hellas (FORTH), P.O. Box 1527, Iraklio, Crete 71110, Greece**

(Dated: June 2005)

We calculate the low temperature one-particle scattering rate and the specific heat in a weakly disordered metal close to a quantum critical point. To lowest order in the fluctuation potential, we obtain typical Fermi-liquid results proportional to T^2 and T respectively, with prefactors which diverge as a power law of the control parameter upon approaching the critical point. The Kadowaki-Woods ratio is shown to be independent of the control parameter only for the case of 3-D FM fluctuations. Our work is relevant for experiments on CeCoIn₅ and Sr₃Ru₂O₇.

We calculate the one-particle scattering rate as a function of temperature $T \rightarrow 0$. We use the relation

$$\text{Im}\Sigma^R(k, \epsilon) = \sum_q \int_{-\infty}^{\infty} d\omega \text{Im}G^R(q, \epsilon - \omega) \text{Im}V^R(k - q, \omega) \{ \coth(\omega/2T) + \tanh((\epsilon - \omega)/2T) \} , \quad (1)$$

to calculate the scattering rate, which equals twice $\text{Im}\Sigma^R$ - c.f. fig 1(a). This is to lowest order in the magnetic fluctuation potential. Here

$$G(k, \epsilon) = \frac{1}{\epsilon - \epsilon_k + i \text{sgn}(\epsilon)/2\tau} \quad (2)$$

with τ being the impurity momentum relaxation time.

The dominant electron-electron interaction is assumed to be the fluctuation propagator^{1,2} (finite q_0 was considered in³)

$$V(q, \omega) = \frac{g}{-i\omega/(Dq^2) + (q - q_0)^2\xi^2 + a} , \quad (3)$$

with a measuring the distance from the quantum critical point (QCP). The control parameter a depends on e.g. the magnetic field H , as in the systems of interest mentioned below like $a = h^s$ - typically $s = 2/3$ - $h = (H - H_c)/H_c$. $q_0=0$ corresponds to ferromagnetic (FM) fluctuations.

Carrying-out this calculation yields in 3-D

$$\text{Im}\Sigma^R(k_F, \epsilon_F, T) = \text{const.} T^2 f_3(a, q_0) , \quad (4)$$

with

$$f_3(a, q_0) = \frac{1}{2a\xi^2 q_0} \left\{ \frac{1}{q_{max} - q_0} + \frac{1}{q_0} \right\} , \quad q_0 = \text{finite} , \quad (5)$$

$$= \frac{1}{a^2} , \quad q_0 = 0 . \quad (6)$$

In 2-D we have

$$\text{Im}\Sigma^R(k_F, \epsilon_F, T) = \text{const.} T^2 f_2(a, q_0) \left\{ \ln \left(\frac{T_o}{T} \right) - 1 + \frac{3T}{T_o} + O(T^2) \right\} , \quad T_o = \frac{a^2 D'}{\xi^2} . \quad (7)$$

It turns out that $f_2(a, q_0) = f_3(a, q_0)$, so the prefactor dependence on a is the same as in 3-D.

Taking the resistivity proportional to the scattering rate, these results compare very favorably with data on CeCoIn₅⁴ and Sr₃Ru₂O₇⁵, where a typical Fermi-liquid (FL) T -dependence is seen in the resistivity $\rho(T) = \rho_0 + A(H)T^2$, with $A(H)$ diverging around the transition, i.e. around $a = 0$, as in our result. In CeCoIn₅ it was found that $A(H) \propto a^{-2}$, as in the case $q_0 = 0$ above. The same result was obtained for the thermal conductivity⁴, which is easily understood in the frame of our calculation: conduction electrons carry both charge and heat, while interacting via $V(q, \omega)$ (only *small* energy transfer is involved with this $V(q, \omega)$).

It should be possible to probe the scattering rate through angle-resolved photoemission (ARPES) experiments.

We calculate the specific heat $C(T) = -T\partial^2 F(T)/\partial T^2 = \gamma T$ through the free energy $F(T)$ - c.f. fig 1(b). In 3-D we obtain for all q_0

$$\gamma = \text{const.}/(a \xi^2) . \quad (8)$$

In 2-D we have for all q_0

$$\gamma = \text{const.}/\xi^2 , \quad (9)$$

i.e. there is no diverging prefactor.

The Kadowaki-Woods ratio is

$$\frac{A(a)}{\gamma^2(a)} = c_1 \xi^4 , \quad 3 - D , \quad q_0 = 0 , \quad (10)$$

$$= c_2 a \xi^2 , \quad 3 - D , \quad q_0 = \text{finite} , \quad (11)$$

$$= c_3 \xi^4 / a^2 , \quad 2 - D , \quad q_0 = 0 , \quad (12)$$

$$= c_4 \xi^2 / a , \quad 2 - D , \quad q_0 = \text{finite} , \quad (13)$$

$c_{1-4} = \text{const.}$ It is independent of a only in 3-D for $q_0 = 0$, while some simple power law arises in the other cases considered. An a -independent ratio was observed in CeCoIn₅⁶, though in a more restricted range of H than the scaling of the coefficient $A(H)$.

We also consider an interaction with peaks at specific wavevectors $\vec{q}_{0i}$ ("AF" case)

$$V_*(q, \omega) = \sum_{i=1}^{n_d} \frac{g}{-i\omega/(Dq^2) + (\vec{q} - \vec{q}_{0i})^2 \xi^2 + a} . \quad (14)$$

In 2-D for tetragonal symmetry $n_d = 4$ and in 3-D for cubic symmetry $n_d = 6$. For small a , in 2-D and 3-D the potential $V_*(q, \omega)$ gives the same scaling of the prefactors as for the case 3-D $q_0 = \text{finite}$ above.

In all, a consistent Fermi liquid (FL) description seems to emerge from these calculations. The renormalization of the fermions due to $V(q, \omega)$ leads back to the FL fixed point in a low- T part of the phase diagram - e.g. c.f. fig 1 of ref.⁴(case $n = 2$). The effective mass of the fermions is not renormalized to first order in $V(q, \omega)$ - c.f. the match between the one particle scattering rate and the experimental resistivity, connected through $\rho = m_* \tau^{-1}/(ne^2)$.

* e-mail : kast@iesl.forth.gr

¹ J. Hertz, Phys. Rev. B **14**, 1165 (1976).


² Y.B. Kim and A.J. Millis, Phys. Rev. B **67**, 085102 (2003).

³ A.V. Chubukov, V.M. Galitski and V.M. Yakovenko, Phys. Rev. Lett. **94**, 046404 (2005).


⁴ J. Paglione et al., cond-mat/0405157.

⁵ S.A. Grigera et al., Science **294**, 329 (2001).

⁶ A. Bianchi et al., Phys. Rev. Lett. **91**, 257001 (2003).



(a)



(b)

Fig. 1

FIG. 1: Diagrams (a) for the self-energy and (b) for the free energy. The continuous lines are the fermions and the dashed line is the fluctuation mediated interaction $V(q, \omega)$.